

Applicants: Bradley et al.  
U.S.S.N. 09/839,658

## IN THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

### Listing of Claims:

1. (currently amended) A method for generating a molecular profile of genomic DNA by hybridization of a genomic DNA target to a plurality of immobilized nucleic acid probes, wherein the plurality is a collection of clones that represent all ~~or part~~ of a chromosome or a genome of an organism, the method comprising:
  - (a) providing the plurality of nucleic acid probes comprising a plurality of immobilized nucleic acid segments in an array with each probe at a known location, wherein each probe is a member of a genomic library cloned in a vector, and each probe is the vector having a cloned nucleic acid insert greater than about 50 kilobases, wherein the plurality of probes represents all ~~or part~~ of the chromosome or the genome;
  - (b) contacting the immobilized probes with a sample of target nucleic acid comprising fragments of genomic nucleic acid labeled with a detectable moiety, wherein each labeled fragment consists of a length smaller than about 200 bases, and the contacting is under conditions allowing hybridization of the target nucleic acid to the probe nucleic acid; and
  - (c) observing an amount and location of labeled genomic nucleic acid hybridized to each immobilized probe, to detect regions of amplification, or deletion ~~and unique sequences~~ in the sample, wherein positional information of clones on the arrays and chromosomes is correlated, thereby generating a molecular profile of the chromosome or genome of the sample genomic nucleic acid.
2. (original) The method of claim 1, wherein each labeled fragment consists of a length no more than about 150 bases.

Applicants: Bradley et al.  
U.S.S.N. 09/839,658

3. (original) The method of claim 2, wherein each labeled fragment consists of a length no more than about 100 bases.
4. (original) The method of claim 3, wherein each labeled fragment consists of a length no more than about 50 bases.
5. (original) The method of claim 4, wherein each labeled fragment consists of a length no more than about 30 bases.
6. (previously presented) The method of claim 2, wherein each labeled fragment consists of a length between about 30 bases and about 150 bases.
7. (previously presented) The method of claim 1, wherein the sample of target genomic nucleic acid is prepared using a procedure selected from the group consisting of random priming, nick translation, and amplification, of a sample of genomic nucleic acid to generate segments of target genomic nucleic acid; followed by a step comprising fragmentation or enzymatic digestion, or both, of the segments to generate a sample of target genomic nucleic acid consisting of sizes smaller than about 200 bases.
8. (previously presented) The method of claim 7, wherein the random priming, nick translation, or amplification, of the sample of genomic nucleic acid to generate segments of target genomic nucleic acid incorporates detectably labeled base pairs into the segments.
9. (previously presented) The method of claim 8, wherein the detectable label comprises Cy3™ or Cy5™.
10. (previously presented) The method of claim 1, further comprising prior to step (b), fragmenting the sample of target genomic nucleic acid to sizes smaller than about 200 bases by DNase enzyme digestion.

Applicants: Bradley et al.  
U.S.S.N. 09/839,658

11. (previously presented) The method of claim 1, further comprising prior to step (b), fragmenting the sample of target genomic nucleic acid to sizes smaller than about 200 bases by applying shearing forces sufficient to fragment genomic DNA followed by DNase enzyme digestion of the sheared DNA.
12. (original) The method of claim 1, wherein the conditions allowing hybridization of the target nucleic acid to the probe nucleic acid comprise stringent hybridization conditions.
13. (original) The method of claim 12, wherein the stringent hybridization conditions comprise a temperature of about 60°C to about 65°C.
14. (original) The method of claim 1, wherein the target nucleic acid consists essentially of DNA derived from a human.
- 15-16. (canceled)
17. (currently amended) The method of claim 1 ~~15 or 16~~, wherein the chromosome or genome is derived from a human.
- 18-66. (canceled)
67. (previously presented) The method of claim 1, wherein the sample of target genomic nucleic acid comprises at least one chromosome.
68. (previously presented) The method of claim 1, wherein the sample of target genomic nucleic acid comprises a complete genome.